



NW Center for Colorectal Health, LLC
9155 SW Barnes Rd, Suite 231
Portland, OR 97225
Phone 503.216-5380
Fax 503.216-5399

Scott M. Browning, MD, FACS, FASCRS
Megan M. Cavanaugh, MD, FACS, FASCRS
Robert C. Amajoyi, MD
Physicians and Surgeons
www.nwcch.com

This diet consists of clear liquids with little nutritional value, and should be followed only for a short period of time. One or two days of this diet does not represent a health risk. A clear liquid is defined as any liquid you can shine a light through, that is not foggy or cloudy and is not colored red or purple.

BEVERAGES: Apple juice, **white** grape juice, **white** cranberry juice, strained lemonade or limeade, tea (no milk, cream or nondairy creamer), water, carbonated and non-carbonated soft drinks, Kool-Aid and Gatorade.

DESSERTS: Italian ices, ice-popsicles, Jell-O **without** whipped topping or fruit, and hard candy.

SOUP: Clear broth or bouillon without noodles, vegetables, etc.

RESTRICTIONS: AVOID any **red or purple colored** clear liquids, including gelatins, beverages and ices. .

NON-INSULIN DEPENDANT DIABETCS: GENERAL GUIDELINES & INSTRUCTIONS

- ◆ We encourage you to contact your PCP or endocrinologist to discuss proper glucose regulation during your prep.
- ◆ Your blood glucose and vital signs will be monitored while at the facility.
- ◆ Bring your oral hypoglycemic medication to the procedure. You can take it after the procedure is over if necessary.
- ◆ Oral hypoglycemic medications generally do not need adjustments and patients may take their oral medication as directed up until the day of the procedure.
 - For AM procedures: You should hold the AM oral medication dose and take it after the procedure is over.
 - For PM procedures: you may take clear liquids in the AM up to 4 hours before the procedure. You may suck on clear hard candies or chew on sugared gum up to the procedure time for low blood sugar.
- ◆ Patients taking **insulin** should only **take half (1/2) the usual scheduled dose** before the procedure. Please call your prescribing physician with any questions regarding your medications.

GENERAL HINTS & REMINDERS FOR COLONOSCOPY

- ◆ Consider trying flushable baby wipes to help avoid irritation from wiping with toilet paper.
- ◆ You may use Vaseline or any water soluble lubricants for relief from any rectal irritation. **DO NOT** use Desitin ointment.
- ◆ The laxative will vary in how quickly it works from patient to patient. You may experience diarrhea within an hour, or you may have to wait several hours.
- ◆ Be prepared to describe your last few bowel movements. You will be asked about the color and appearance to make sure the preparation has worked.
- ◆ Rinsing your mouth and gargling with mouthwash can eliminate any aftertaste between doses of preparation liquid.
- ◆ If you feel nauseated or bloated while taking the prep, you may stop the prep for 30-45 minutes one time and then resume.
- ◆ If you use reading glasses, please bring them with you as you will need to read and sign a consent form and discharge sheet.